

Essential Knowledge Milestones	Teaching Points
<ul style="list-style-type: none"> <li>• be able to find the binomial expansion of <math>(1 - x)^{-1}</math> for rational values of <math>n</math> and <math> x  &lt; 1</math>;</li> <li>• be able to find the binomial expansion of <math>(1 + x)^n</math> for rational values of <math>n</math> and <math> x  &lt; 1</math>;</li> <li>• be able to find the binomial expansion of <math>(1 + bx)^n</math> for rational values of <math>n</math> and <math> x  &lt; \frac{1}{ b }</math>;</li> <li>• be able to find the binomial expansion of <math>(a + x)^n</math> for rational values of <math>n</math> and <math> x  &lt; a</math>;</li> <li>• be able to find the binomial expansion of <math>(a + bx)^n</math> for rational values of <math>n</math> and <math> \frac{bx}{a}  &lt; 1</math>;</li> <li>• know how to use the binomial theorem to find approximations (including roots).</li> <li>• be able to use partial fractions to write a rational function as a series expansion.</li> </ul>	<ul style="list-style-type: none"> <li>• Begin by reviewing the expansion of <math>(a + b)^n</math> when <math>n</math> is a positive integer.</li> <li>• Ask students to expand <math>(1 + x)^4</math> and then try <math>(1 + x)^{-2}</math>. Why does it fail to work? Which coefficient calculation breaks down?</li> <li>• Explain how the binomial theorem allows us to expand any power. (Explain the reasoning behind the factorial notation using the explanation in the <i>Reasoning and problem solving</i> section below.)</li> <li>• Consider why the expansions are infinite when the power is <i>not</i> a positive integer. How far do we need to expand and to which term? (For example, up to and including coefficients of <math>x^3</math>.)</li> <li>• Take care to show the precision needed when dealing with negative calculations by demonstrating examples such as <math>(1 - 2x)^{-\frac{1}{2}}</math>.</li> <li>• If we expanded <math>(1 + x)^{\frac{1}{2}}</math> then substituted <math>x = -0.1</math>, we would be effectively finding the square root of 0.9. Ask students to use a calculator to find an accurate value for <math>\sqrt{0.9}</math>. How many terms of the expansion would we need to substitute into in order to get a 4 decimal place version of the accurate value?</li> <li>• What happens when we substitute <math>x = 3</math>? Does this find the square root of 4?</li> <li>• Explain that if we raise a number <math>&gt; 1</math> to a positive power, it 'grows' and diverges out of control. This means that the value of <math>x</math> must be such that <math>-1 &lt; x &lt; 1</math> or <math> x  &lt; 1</math> in order to use the expansion of <math>(1 + x)^n</math>. The validity of the expansion is dependent upon the value of <math>x</math> we substitute into the terms.</li> <li>• Cover examples that build-up the expansions listed in the objectives above, ending with <math>(a + bx)^n</math> for rational values of <math>n</math> and valid for <math> \frac{bx}{a}  &lt; 1</math>.</li> <li>• Introduce the concept of expansions of expressions which start with <math>a</math> rather than 1. Begin by showing that if we have <math>(2 + x)</math> and if we want to make this start with a 1 in the bracket, we must take out the factor of 2, giving <math>2(1 + \frac{x}{2})</math>.</li> <li>• Now show for example, that <math>2(1 + 4)</math> gives the same result as <math>(2 + 8)</math> if we multiplied this out, but that if the bracket were squared the result would not be the same i.e. <math>2(1 + 4)^2 \neq (4 + 8)^2</math>.</li> <li>• However, <math>2^2(1 + 4)^2 = (4 + 8)^2</math>, so we need to raise the factor to the same power of the bracket and <math>(a + bx)^n = a^n(1 + \frac{bx}{a})^n</math>.</li> <li>• Need to break-up a rational function into two or more partial fractions.</li> <li>• If we consider the 'complicated' fraction below, it needs to be simplified into two simpler fractions each of which only involve a single algebraic bracket.</li> <li>• <math display="block">\frac{2x^2+5x-10}{(x-1)(x+2)} \equiv A + \frac{B}{x-1} + \frac{C}{x+2}</math></li> </ul>
Assumed Prior Knowledge/ Links / Interleaving	
<ul style="list-style-type: none"> <li>• GCSE: Simplifying algebraic fractions</li> <li>• The Binomial Expansion: this looks at the finite expansion of <math>(a + bx)^n</math>, where <math>n</math> is a positive integer.</li> <li>• Infinite geometric series: <math>1 + x + x^2 + x^3 + x^4 + \dots = (1 - x)^{-1}</math>; geometric series start from the left hand side, in this unit we start with the right hand side.</li> <li>• Polynomials: In this unit we see that over a certain domain all functions of the form <math>(1 + x)^n</math> can be approximated by polynomials.</li> <li>• Integration: partial fractions transforms some expressions into a form that allows us to integrate them</li> </ul>	

	<ul style="list-style-type: none"> <li>We can now rewrite each term as a binomial series. (It is important to demonstrate that the <math>\frac{B}{x-1}</math> term will become <math>B(x-1)^{-1}</math>.)</li> <li>Particular care needs to be taken when working with brackets that don't start with 1, and also when multiplying out all the terms to arrive at the final simplified series (up to and including the power required).</li> </ul>	
<b>Potential Barriers to Access/Misconceptions</b>	<b>Opportunities for Reasoning/Problem Solving/Proofs</b>	
<ul style="list-style-type: none"> <li>When expanding <math>(1 + 4x)^{\frac{1}{2}}</math> most students got the first two terms of the expansion correct, but often there was a mistake in the <math>x^2</math> term, with <math>4x</math> becoming just <math>x</math> being the common error. Some students made arithmetic errors with <math>4^2</math>, by failing to actually square the 4, and others failed to simplify the binomial coefficient correctly.</li> <li>When expanding an expression of the form <math>(a + x)^n</math> a common error is to write this as <math>a(1 + \frac{x}{a})^n</math> rather than <math>a^n(1 + \frac{x}{a})^n</math>.</li> <li>Other errors include algebraic errors when combining two expansions, doing more work than is necessary when, for example, only terms up to <math>x^2</math> are required, including the equality in the expression for the range of valid values for <math>x</math> and lack of understanding when using the modulus symbol (writing expressions such as <math> x  &lt; -4</math>).</li> <li>misinterpreting <math>(a\sqrt{b})^2</math> as <math>(a + \sqrt{b})^2</math>;</li> <li>Expanding <math>(a + bx)^n</math> as if <math>a</math> had the value 1.</li> <li>Mistakes over signs in expanding, for example, <math>(1 - 3x)^{-2}</math></li> <li>Missing out partial fractions when there is a repeated factor</li> <li>Omitting the 'validity term' or stating it incorrectly.</li> <li>Able to write <math>(x - 1)^{-1}</math> as <math>-(1 - x)^{-1}</math> and the resulting expansions were incorrect in the majority of cases, both <math>1 + x - x^2</math> and <math>1 - x - x^2</math> being common errors.</li> </ul>	<ul style="list-style-type: none"> <li>Using the binomial expansion of <math>(1 - \frac{1}{2})^{-\frac{1}{2}}</math>, prove that <math>\sqrt{2} &gt; \frac{32}{23}</math>. Can you find a better rational approximation for <math>\sqrt{2}</math>?</li> <li>Show how <math>{}^nC_r</math> will only work on a calculator for positive integer values of <math>n</math> (as was done in Unit 3 of Pure).</li> <li>However, we can instead use the definition and formula for selections (shown here for 'choose 2 from <math>n</math> different objects').</li> <li> <math display="block">{}^nC_2 = \frac{n!}{2!(n-2)!}</math> <math display="block">= \frac{n(n-1)(\cancel{n-2})(\cancel{n-3})\dots}{2!(\cancel{n-2})(\cancel{n-3})\dots}</math> <math display="block">= \frac{n(n-1)}{2!}</math> </li> <li>This formula works for <i>all</i> values of <math>n</math> and follows the pattern of the binomial theorem as stated in the formula book.</li> <li>You will need to assess all the separate validities for the individual binomial terms to declare the validity for the final series.</li> <li>Include examples in which one of the terms is not a binomial and just multiplies without expansion. For example, <math>(2x - 1)(1 + 3x)^{-2}</math>.</li> </ul>	
	<b>Questions &amp; Prompts</b>	
	<ul style="list-style-type: none"> <li>Explain how the binomial theorem can be used to find approximations of square roots.</li> <li>Give me an example of a binomial expansion where the expansion is valid only for <math> x  &lt; \frac{1}{3}</math>.</li> <li>Make up three questions that show you understand different applications of partial fractions.</li> </ul>	
<b>Key Mathematical Vocabulary</b>	Binomial, expansion, theorem, integer, rational, power, index, coefficient, validity, modulus, factorial, ${}^nC_r$ , combinations, Pascal's triangle, partial fractions, approximation, converges, diverges, root.	
<b>Personal Development</b>	<b>Notes</b>	<b>Resources</b>
Pupils are taught that they must show 'ambition' in their assignments by attempting to meet all criteria in their homework, with honesty with regards what support they have had to complete the assignment		